

## Referencias bibliográficas

- Al-Delaimy, W. K., Ferrari, P., Slimani, N., Pala, V., Johansson, I., Nilsson, S., Mattisson, I., Wirfalt, E., Galasso, R., Palli, D., Vineis, P., Tumino, R., Dorransoro, M., Pera, G., Ocké, M. C., Bueno-de-Mesquita, H. B., Overvad, K., Chirlaque, M., Trichopoulou, A., ... Riboli, E. (2005). Plasma carotenoids as biomarkers of intake of fruits and vegetables: individual-level correlations in the European Prospective Investigation into Cancer and Nutrition (EPIC). *European Journal of Clinical Nutrition*, 59(12), 1387–1396. <https://doi.org/10.1038/sj.ejcn.1602252>
- Allen, N. E., Grace, P. B., Ginn, A., Travis, R. C., Roddam, A. W., Appleby, P. N., & Key, T. (2008). Phytanic acid: measurement of plasma concentrations by gas–liquid chromatography–mass spectrometry analysis and associations with diet and other plasma fatty acids. *British Journal of Nutrition*, 99(3), 653–659. <https://doi.org/10.1017/S000711450782407X>
- Arsenault, L. N., Matthan, N., Scott, T. M., Dallal, G., Lichtenstein, A. H., Folstein, M. F., Rosenberg, I., & Tucker, K. L. (2009). Validity of Estimated Dietary Eicosapentaenoic Acid and Docosahexaenoic Acid Intakes Determined by Interviewer-Administered Food Frequency Questionnaire Among Older Adults With Mild-to-Moderate Cognitive Impairment or Dementia. *American Journal of Epidemiology*, 170(1), 95–103. <https://doi.org/10.1093/aje/kwp089>
- Baldrick, F. R., Woodside, J. v., Elborn, J. S., Young, I. S., & McKinley, M. C. (2011). Biomarkers of Fruit and Vegetable Intake in Human Intervention Studies: A Systematic Review. *Critical Reviews in Food Science and Nutrition*, 51(9), 795–815. <https://doi.org/10.1080/10408398.2010.482217>
- Bictash, M., Ebbels, T. M., Chan, Q., Loo, R. L., Yap, I. K. S., Brown, I. J., de Iorio, M., Daviglus, M. L., Holmes, E., Stamler, J., Nicholson, J. K., & Elliott, P. (2010). Opening up the “Black Box”: Metabolic phenotyping and metabolome-wide association studies in epidemiology. *Journal of Clinical Epidemiology*, 63(9), 970–979. <https://doi.org/10.1016/j.jclinepi.2009.10.001>
- Brantsæter, A. L., Haugen, M., Julshamn, K., Alexander, J., & Meltzer, H. M. (2009). Evaluation of urinary iodine excretion as a biomarker for intake of milk and dairy products in pregnant women in the Norwegian Mother and Child Cohort Study (MoBa). *European Journal of Clinical Nutrition*, 63(3), 347–354. <https://doi.org/10.1038/sj.ejcn.1602952>
- Freedman, L. S., Potischman, N., Kipnis, V., Midthune, D., Schatzkin, A., Thompson, F. E., Troiano, R. P., Prentice, R., Patterson, R., Carroll, R., & Subar, A. F. (2006). A comparison of two dietary instruments for evaluating the fat–breast cancer relationship. *International Journal of Epidemiology*, 35(4), 1011–1021. <https://doi.org/10.1093/ije/dyl085>
- LeVatte, M., Keshteli, A. H., Zarei, P., & Wishart, D. S. (2022). Applications of Metabolomics to Precision Nutrition. *Lifestyle Genomics*, 15(1), 1–9. <https://doi.org/10.1159/000518489>
- Lloyd, A. J., Beckmann, M., Favé, G., Mathers, J. C., & Draper, J. (2011). Proline betaine and its biotransformation products in fasting urine samples are potential biomarkers of habitual citrus fruit consumption. *British Journal of Nutrition*, 106(6), 812–824. <https://doi.org/10.1017/S0007114511001164>
- Mennen, L. I., Sapinho, D., Ito, H., Bertrais, S., Galan, P., Hercberg, S., & Scalbert, A. (2006). Urinary flavonoids and phenolic acids as biomarkers of intake for polyphenol-rich foods. *British Journal of Nutrition*, 96(01), 191. <https://doi.org/10.1079/BJN20061808>
- Pirro, V., Valente, V., Oliveri, P., de Bernardis, A., Salomone, A., & Vincenti, M. (2011). Chemometric evaluation of nine alcohol biomarkers in a large population of clinically-classified subjects: pre-eminence of ethyl glucuronide concentration in hair for confirmatory classification. *Analytical and Bioanalytical Chemistry*, 401(7), 2153–2164. <https://doi.org/10.1007/s00216-011-5314-7>
- Ross, A. B., Bourgeois, A., Macharia, H. N., Kochhar, S., Jebb, S. A., Brownlee, I. A., & Seal, C. J. (2012). Plasma alkylresorcinols as a biomarker of whole-grain food consumption in a large population: results from the WHOLEheart Intervention Study. *The American Journal of Clinical Nutrition*, 95(1), 204–211. <https://doi.org/10.3945/ajcn.110.008508>
- Scalbert, A., Brennan, L., Manach, C., Andres-Lacueva, C., Dragsted, L. O., Draper, J., Rappaport, S. M., van der Hoof, J. J., & Wishart, D. S. (2014). The food metabolome: a window over dietary exposure. *The American Journal of Clinical Nutrition*, 99(6), 1286–1308. <https://doi.org/10.3945/ajcn.113.076133>
- Turunen, A. W., Männistö, S., Kiviranta, H., Marniemi, J., Jula, A., Tiittanen, P., Suominen-Taipale, L., Vartiainen, T., & Verkasalo, P. K. (2010). Dioxins, polychlorinated biphenyls, methyl mercury and omega-3 polyunsaturated fatty acids as biomarkers of fish consumption. *European Journal of Clinical Nutrition*, 64(3), 313–323. <https://doi.org/10.1038/ejcn.2009.147>
- Vinayavekhin, N., & Saghatelian, A. (2010). Untargeted Metabolomics. In *Current Protocols in Molecular Biology*. John Wiley & Sons, Inc. <https://doi.org/10.1002/0471142727.mb3001s90>